


Team 1

Plate 1a Nitrate and Conductivity Record, GISP2-H-Part I Section 1

Dreschhoff and Zeller  
TER-QUA Proceedings, V.




Team 2

Plate 1a Nitrate and Conductivity Record, GISP2-H-Part I Section 2

1994.

DALTON MINIMUM →




Team 3

Plate 1b Nitrate and Conductivity Record, GISP2-H-Part II Section 1

Dreschhoff and Zeller  
TER-QUA Proceedings, Volume 2, 1994.

DALTON MINIMUM ←

MAUNDER MINIMUM →



Team 4

Plate 1b Nitrate and Conductivity Record, GISP2-H-Part II Section 2

1994.

MAUNDER MINIMUM ←

Plates Ia and Ib

Micro-resolution records of simultaneous nitrate concentrations in relative units (blue curve) and liquid conductivity (orange curve) from the Central Greenland Ice Sheet. The x-axis is proportional to the true depth below the surface (122m) and has not been adjusted for compaction. Nitrate background displays a prominent seasonal cycle. The nitrate anomalies are apparently caused in most cases by individual solar particle events (dates have been marked in blue). Black arrows at the bottom of the graph indicate times of sunspot maxima (up) and sunspot minima (down). The numbers at the top of the graph indicate the approximate summer peaks of the years of the most recent maximum in solar activity. Asterisks next to major peaks indicate that these anomalies are part of the ongoing evaluation programs. Areas of gray shading represent portions of the core resampled for confirmation. Absorbance units ( $100 \times 10^{-4}$ ) convert to  $20.4 \mu\text{g/l NO}_3\text{-N}$  or  $90.3 \text{ ng/g NO}_3$ . Anomalous conductivity peaks indicate the years of specific volcanic eruptions.

